

ENEMA - mineral oil enema

Amerisource Bergen Drug Corporation

DRUG FACTS

ACTIVE INGREDIENT

Mineral Oil, 100%

PURPOSE

Lubricant Laxative

USE

- For relief of occasional constipation. This product generally produced bowel movement in 2 to 15 minutes.

WARNINGS

Using more than one enema in 24 hours can be harmful.

Do not use

- If you have abdominal pain, nausea, or vomiting, unless directed by a doctor.
- Longer than one week unless directed by a doctor.
- More than one enema per day unless directed by a doctor.

Consult a doctor before use

- if you have noticed a sudden change in bowel habits that persist over a period of 2 weeks
- if you have rectal bleeding or failure to have a bowel movement after use of a laxative. This may indicate a serious condition.
- if you are pregnant or nursing a baby

Keep out of reach of children

In case of accidental injection or overdose, get medical help or contact a Poison Control Center right away.

DIRECTIONS

Daily Rectal Dosage

Adults and children 12 years and older	One bottle (4 oz.) 120 mL delivered dose
Children 2 years to 11 years old	1/2 Adult dose (2 oz.) 60 mL delivered dose
Children under 2 years of age	CONSULT A DOCTOR

Dosage

- Adults and children 12 years of age and older: rectal enema dosage is 120 milliliters in a single daily dose.
- Children 2 to 11 years of age: rectal enema dosage is 60 milliliters in a single daily dose, 1/2 of a bottle. (Children's dose of 60 milliliters can be measured by removing mineral oil from the bottle until volume is just under the words "Mineral Oil" on the front face of the bottle).
- Children under 2 years of age:
Consult a doctor.

Directions for inserting enema

CAUTION: Remove orange protective shield before inserting. To do so, hold bottle upright with one hand and pull gently to remove.

Positioning:

Left side position: Lie on left side with right knee bent and arms resting comfortably.

Knee chest position: Kneel, lower head and chest forward until left side of the face is resting on surface with arms resting comfortably.

- With steady pressure, gently insert enema tip into rectum with a side to side movement with tip pointing toward navel.
- Stop using if tip is hard to insert. Forcing tip into the rectum can cause injury (especially if you have hemorrhoids). If enema tip causes rectal bleeding or pain, get immediate medical care.
- Squeeze bottle until nearly all the liquid is released. Remove from rectum. Maintain position until urge to evacuate is strong (usually 2 to 15 minutes).

OTHER INFORMATION

- This product does not contain Dry Natural Rubber or Natural Rubber Latex.
- Store at room temperature 15° - 30° C (59° - 86° F)
- Enema tip coated with petrolatum for ease of insertion

PACKAGE INFORMATION

NDC# 24385-360-28

GOOD NEIGHBOR PHARMACY®

Ready to Use
Mineral Oil, USP
ENEMA

LUBRICANT LAXATIVE FOR THE RELIEF OF OCCASIONAL CONSTIPATION

* COMPARE TO THE ACTIVE INGREDIENT IN FLEET®

Single use
4.5 FL. OZ. (133 ml)

- Orange protective shield prevents contamination
- Lubricated tip assures ease of insertion

- One way safety valve prevents reflux
- Easy to grip and squeeze bottle

CARTON SEALED FOR SAFETY. DO NOT USE IF GLUE SEAL AT EITHER END IS BROKEN OR HAS BEEN DISTURBED.

*This product is not manufactured or distributed by C.B. Fleet Co., owner of the registered trademark Fleet®.

Distributed by
Amerisource Bergen
Valley Forge, PA 19482-0959

ENEMA
Mineral Oil, USP
Ready To Use



Drug Facts (continued)

Directions for inserting enema

CAUTION: Remove orange protective shield before inserting. To do so, hold bottle upright with one hand and pull gently to remove.

Left Side Position: Lie on left side with right knee bent and arms resting comfortably.



Knee Chest Position: Kneel, lower head and chest forward until left side of the face is resting on surface with arms resting.



- With steady pressure, gently insert enema tip into rectum with a side to side movement with tip pointing toward navel.
- Stop using if tip is hard to insert. Forcing tip into the rectum can cause injury (especially if you have hemorrhoids); if enema tip causes rectal bleeding or pain, get immediate medical care.
- Squeeze bottle until nearly all the liquid is released. It is not necessary to empty the bottle completely as it contains more liquid than needed. Remove from rectum. Maintain position until urge to evacuate is

Other information

- This product does not contain Dry Natural Rubber or Natural Rubber Latex.
- Store at room temperature: 15° - 30° C (59° - 86° F).
- Enema tip coated with petroleum for ease of insertion.

*This product is not manufactured or distributed by C.B. Fleet Co. owner of



Distributed By
AmerisourceBergen
Valley Forge, PA. 19482-0959

Visit us at
www.myGNP.com



Ready To Use
Mineral Oil, USP
ENEMA

- Orange protective shield prevents contamination
- Lubricated tip assures ease of insertion

- One way safety valve prevents reflux
- Easy to grip and squeeze bottle



NDC# 24385-360-28



Ready To Use
Mineral Oil, USP
ENEMA

LUBRICANT LAXATIVE
FOR THE RELIEF OF
OCCASIONAL

*COMPARE TO
THE ACTIVE
INGREDIENT
IN FLEET®



Single use
4.5 FL. OZ. (133 mL)

Rev. A

100% UPC
F.P.O.



ABCP 245-530
4483772

Drug Facts

Active ingredient Mineral Oil, USP 100% **Purpose** Lubricant Laxative

Use
Rectal lubricant/laxative for relief of occasional constipation. This product generally produces bowel movement in 7 to 15 minutes.

Warnings
Using more than one enema in 24 hours can be harmful.

Do not use

- if you have abdominal pain, nausea, or vomiting, unless directed by a doctor.
- longer than one week unless directed by a doctor.
- more than one enema per day unless directed by a doctor.

Consult a doctor prior to use

- if you have noticed a sudden change in bowel habits that persists over a period of 2 weeks.
- if you have rectal bleeding or failure to have a bowel movement after use of a laxative. This may indicate a serious condition.
- if you are pregnant or nursing a baby.

Keep out of reach of children. In case of accidental ingestion or overdose, get medical help or contact a Poison Control Center right away.

Directions

Daily Rectal Dosage	
Adults and children 12 years and older	One bottle (4 oz.) 133 mL, delivered dose
Children 2 years to 11 years old	1/2 Adult Dose (2 oz.) 60 mL, delivered dose
Children under 2 years of age	CONSULT A DOCTOR

- Dosage**
- **Adults and children 12 years of age and older:** Rectal enema dosage is 133 mL in a single daily dose.
 - **Children 2 to 11 years of age:** Rectal enema dosage is 60 mL in a single daily dose, 1/2 of 1 bottle. (Children's dose of 60 mL can be measured by removing mineral oil from the bottle until volume is just below the words "Mineral Oil" on the front face of the bottle).
 - **Children under 2 years of age:**

CARTON SEALED FOR SAFETY. DO NOT USE IF GLUE SEAL AT EITHER END IS BROKEN OR HAS

ENEMA

mineral oil enema

Product Information

Product Type	HUMAN OTC DRUG	NDC Product Code (Source)	24385-360
Route of Administration	RECTAL	DEA Schedule	

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
mineral oil (mineral oil)	mineral oil	1 mL in 1 mL

Inactive Ingredients

Ingredient Name	Strength
No Inactive Ingredients Found	

Product Characteristics

Color	Score
Shape	Size
Flavor	Imprint Code

Contains

Packaging

# NDC	Package Description	Multilevel Packaging
1 24385-360-28	1 BOTTLE In 1 BOX	contains a BOTTLE
1	133 mL In 1 BOTTLE	This package is contained within the BOX (24385-360-28)

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part334	02/13/2004	03/01/2010

Labeler - Amerisource Bergen Drug Corporation (007914906)

Revised: 02/2010 Amerisource Bergen Drug Corporation